

**QOF Indicators 2006/07**

<b>Indicator Abbreviation</b>	<b>Full indicator description</b>
ATRIAL FIBRILLATION 2	The percentage of patients with atrial fibrillation diagnosed from 1st April 2006 with ECG or specialist confirmed diagnosis
ATRIAL FIBRILLATION 3	The percentage of patients with atrial fibrillation who are currently treated with anti-coagulant drug therapy or an anti-platelet drug therapy
ASTHMA 3	The percentage of patients with asthma between the ages of 14 and 19 in whom there is a record of smoking status in the previous 15 months
ASTHMA 6	The percentage of patients with asthma who have had an asthma review in the previous 15 months
ASTHMA 8	The percentage of patients aged eight and over diagnosed as having asthma from 1st April 2006 with measures of variability or reversibility
HYPERTENSION 4	The percentage of patients with hypertension in whom there is a record of the blood pressure in the previous 9 months
HYPERTENSION 5	The percentage of patients with hypertension in whom the last blood pressure (measured in the previous 9 months) is 150/90 or less
CANCER 3	The percentage of patients with cancer, diagnosed within the last 18 months, who have a patient review recorded as occurring at 6 months after the practice has received confirmation of the diagnosis
CHD 2	The percentage of patients with newly diagnosed angina (diagnosed after 1 April 2003) who are referred for exercise testing and/or specialist assessment
CHD 5	The percentage of patients with coronary heart disease whose notes have a record of blood pressure in the previous 15 months
CHD 6	The percentage of patients with coronary heart disease in whom the last blood pressure reading (measured in the previous 15 months) is 150/90 or less
CHD 7	The percentage of patients with coronary heart disease whose notes have a record of total cholesterol in the previous 15 months
CHD 8	The percentage of patients with coronary heart disease whose last measured total cholesterol (measured in the previous 15 months) is 5 mmol/l or less
CHD 9	The percentage of patients with coronary heart disease with a record in the previous 15 months that aspirin, an alternative anti-platelet therapy, or an anti-coagulant is being taken (unless a contraindication or side-effects are recorded)
CHD 10	The percentage of patients with coronary heart disease who are currently treated with a beta blocker (unless a contraindication or side-effects are recorded)
CHD 11	The percentage of patients with a history of myocardial infarction (diagnosed after 1 April 2003) who are currently treated with an ACE inhibitor or angiotensin II antagonist
CHD 12	The percentage of patients with coronary heart disease who have a record of influenza immunisation in the preceding 1 September to 31 March
CKD 2	The percentage of patients on the CKD register whose notes have a record of blood pressure in the previous 15 months
CKD 3	The percentage of patients on the CKD register in whom the last blood pressure reading, measured in the previous 15 months, is 140/85 or less
CKD 4	The percentage of patients on the CKD register with hypertension, who are treated with an angiotensin converting enzyme inhibitor (ACE-I) or angiotensin receptor blocker (ARB) (unless a contraindication or side effects are recorded)
COPD 8	The percentage of patients with COPD who have had influenza immunisation in the preceding 1 September to 31 March
COPD 9	The percentage of all patients with COPD in whom diagnosis has been confirmed by spirometry including reversibility testing
COPD 10	The percentage of patients with COPD with a record of FeV1 in the previous 15 months
COPD 11	The percentage of patients with COPD receiving inhaled treatment in whom there is a record that inhaler technique has been checked in the previous 15 months
DEMENTIA 2	The percentage of patients diagnosed with dementia whose care has been reviewed in the previous 15 months
DEPRESSION 1	The percentage of patients with diabetes and/or heart disease for whom case finding for depression has been undertaken on one occasion during the previous 15 months using the two standard screening questions
DEPRESSION 2	In those patients with a new diagnosis of depression, recorded between the preceding 1 April and 31st March, the percentage of patients who have had an assessment of severity at the outset of treatment using an assessment tool validated for use in primary care
DIABETES 2	The percentage of patients with diabetes whose notes record BMI in the previous 15 months
DIABETES 5	The percentage of patients with diabetes who have a record of HbA1c or equivalent in the previous 15 months
DIABETES 7	The percentage of patients with diabetes in whom the last HbA1C is 10 or less (or equivalent test/reference range depending on local laboratory) in the previous 15 months
DIABETES 9	The percentage of patients with diabetes with a record of the presence or absence of peripheral pulses in the previous 15 months
DIABETES 10	The percentage of patients with diabetes with a record of neuropathy testing in the previous 15 months
DIABETES 11	The percentage of patients with diabetes who have a record of the blood pressure in the previous 15 months
DIABETES 12	The percentage of patients with diabetes in whom the last blood pressure is 145/85 or less
DIABETES 13	The percentage of patients with diabetes who have a record of micro-albuminuria testing in the previous 15 months (exception reporting for patients with proteinuria)

DIABETES 15	The percentage of patients with diabetes with proteinuria or micro-albuminuria who are treated with ACE inhibitors (or A2 antagonists)
DIABETES 16	The percentage of patients with diabetes who have a record of total cholesterol in the previous 15 months
DIABETES 17	The percentage of patients with diabetes whose last measured total cholesterol within the previous 15 months is 5mmol/l or less
DIABETES 18	The percentage of patients with diabetes who have had influenza immunisation in the preceding 1 September to 31 March
DIABETES 20	The percentage of patients with diabetes in whom the last HbA1C is 7.5 or less (or equivalent test/reference range depending on local laboratory) in the previous 15 months
DIABETES 21	The percentage of patients with diabetes who have a record of retinal screening in the previous 15 months
DIABETES 22	The percentage of patients with diabetes who have a record of estimated glomerular filtration rate (eGFR) or serum creatinine testing in the previous 15 months
EPILEPSY 6	The percentage of patients aged 18 years and over on drug treatment for epilepsy who have a record of seizure frequency in the previous 15 months
EPILEPSY 7	The percentage of patients aged 18 years and over on drug treatment for epilepsy who have a record of medication review involving the patient or carer in the previous 15 months
EPILEPSY 8	The percentage of patients aged 18 years and over on drug treatment for epilepsy who have been seizure free for the last 12 months recorded in the previous 15 months
HEART FAILURE 2	The percentage of patients with a diagnosis of heart failure (diagnosed after the 1st April 2006) which has been confirmed by an echocardiogram or by specialist assessment
HEART FAILURE 3	The percentage of patients with a current diagnosis of heart failure due to LVD who are currently treated with an ACE inhibitor or Angiotensin Receptor Blocker, who can tolerate therapy and for whom there is no contraindication
MENTAL HEALTH 4	The percentage of patients on lithium therapy with a record of serum creatinine and TSH in the previous 15 months
MENTAL HEALTH 5	The percentage of patients on lithium therapy with a record of lithium levels in a therapeutic range within the previous 6 months
MENTAL HEALTH 6	The percentage of patients on the register who have a comprehensive care plan documented in the records agreed between individuals, their family and/or carers as appropriate
MENTAL HEALTH 7	The percentage of patients with schizophrenia, bipolar affective disorder and other psychoses who do not attend the practice for their annual review who are identified and followed up by practice team within 14 days of non-attendance
MENTAL HEALTH 9	The percentage of patients with schizophrenia and bipolar affective disorder and other psychoses with a review recorded in the previous 15 months. In the review there is evidence that the patient has been offered routine health promotion and prevention advice appropriate to their age, gender and health status
SMOKING 1	The percentage of patients with any or any combination of the following conditions: coronary heart disease, stroke or TIA, hypertension, diabetes, COPD or asthma whose notes record smoking status in the previous 15 months. Except those who have never smoked where smoking status need only be recorded once since diagnosis
SMOKING 2	The percentage of patients with any or any combination of the following conditions: coronary heart disease, stroke or TIA, hypertension, diabetes, COPD or asthma who smoke whose notes contain a record that smoking cessation advice or referral to a specialist service, where available, has been offered within the previous 15 months
STROKE 5	The percentage of patients with TIA or stroke who have a record of blood pressure in the notes in the preceding 15 months
STROKE 6	The percentage of patients with a history of TIA or stroke in whom the last blood pressure reading (measured in the previous 15 months) is 150/90 or less
STROKE 7	The percentage of patients with TIA or stroke who have a record of total cholesterol in the previous 15 months
STROKE 8	The percentage of patients with TIA or stroke whose last measured total cholesterol (measured in the previous 15 months) is 5 mmol/l or less
STROKE 10	The percentage of patients with TIA or stroke who have had influenza immunisation in the preceding 1 September to 31 March
STROKE 11	The percentage of new patients with a stroke who have been referred for further investigation
STROKE 12	The percentage of patients with a stroke shown to be non-haemorrhagic, or a history of TIA, who have a record that an anti-platelet agent (aspirin, clopidogrel, dipyridamole or a combination), or an anti-coagulant is being taken (unless a contraindication or side-effects are recorded)
HYPOTHYROIDISM 2	The percentage of patients with hypothyroidism with thyroid function tests recorded in the previous 15 months
CERVICAL SCREENING 1	The percentage of patients aged from 25 to 64 (in Scotland from 21 to 60) whose notes record that a cervical smear has been performed in the last five years